


Mom, where's the MURICIN®?

(mupirocin ointment 2%)

MURICIN Ointment is an FDA veterinary approved, water-removable topical treatment that has bacterial skin infections covered. MURICIN Ointment is indicated for the topical treatment of canine bacterial infections of the skin, including

superficial pyoderma, caused by susceptible strains of *Staphylococcus aureus* and *Staphylococcus intermedius*.



Important information about MURICIN: MURICIN Ointment is not for ophthalmic use. Safety in pregnant or breeding animals has not been determined. Due to a potential hazard for nephrotoxicity due to polyethylene glycol content of the product base, exercise care in use for extensive deep lesions where absorption of large quantities of polyethylene glycol is possible.


For more information call 866-933-2472, e-mail support@dechra.com, or contact your veterinary distributor.

MURICIN[®] Ointment (mupirocin ointment 2%)

ANADA # 200-418. Approved by FDA.


For dermatologic use on dogs

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian. **DESCRIPTION:** Each gram of MURICIN Ointment contains 20 mg of mupirocin in a bland, water-washable ointment base consisting of polyethylene glycol 400 and polyethylene glycol 3350 (polyethylene glycol ointment, NF). Mupirocin is a naturally occurring, broad-spectrum antibiotic. The chemical name is (*E*)-(2*S*,3*R*,4*R*,5*S*)-5-[(2*S*,3*S*,4*S*,5*S*)-2,3-Epoxy-5-hydroxy-4-methylhexyl] tetrahydro-3,4-dihydroxy-β-methyl-2*H*-pyran-2-crotonic acid, ester with 9-hydroxynonanoic acid. The chemical structure is:




CLINICAL PHARMACOLOGY: Mupirocin is a chemical entity produced by fermentation of the organism *Pseudomonas fluorescens*. Mupirocin inhibits bacterial protein synthesis by reversibly and specifically binding to bacterial isoleucyl transfer-RNA synthetase. Due to this mode of action, mupirocin shows no cross-resistance with chloramphenicol, erythromycin, gentamicin, lincomycin, neomycin, novobiocin, penicillin, streptomycin, and tetracycline. Mupirocin is an antimicrobial agent that inhibits the growth of gram-positive and gram-negative bacteria. Bacteria susceptible to the action of mupirocin *in vitro* include the aerobic isolates of *Staphylococcus aureus* (including methicillin-resistant strains and β-lactamase-producing strains), *Staphylococcus intermedius*, *Staphylococcus epidermidis*, other coagulase positive or negative Staphylococci, α-hemolytic Streptococci, β group A Streptococci (including *S. pyogenes*), other β Streptococci (including *S. agalactiae*), group D Streptococci (including *S. faecalis* and *S. faecium*), group Viridans Streptococci, *Streptococcus pneumoniae*, *Corynebacterium hofmannii*, *Bacillus subtilis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Proteus vulgaris*, *Enterobacter cloacae*, *Enterobacter aerogenes*, *Citrobacter freundii*, *Hemophilus influenzae* (including β-lactamase-producing strains), *Neisseria gonorrhoeae* (including β-lactamase-producing strains), *Neisseria meningitidis*, *Branhamella catarrhalis* and *Pasteurella multocida*, and the anaerobic isolates of *Peptostreptococcus anaerobius*, *Clostridium difficile*, and *Clostridium sporogenes*. Clinical


significance of the *in vitro* data is unknown except for susceptible strains of *Staphylococcus aureus* and *Staphylococcus intermedius*. **INDICATIONS FOR USE:** MURICIN Ointment is indicated for the topical treatment of canine bacterial infections of the skin, including superficial pyoderma, caused by susceptible strains of *Staphylococcus aureus* and *Staphylococcus intermedius*. **CONTRAINDICATIONS:** This drug is contraindicated in animals with a history of sensitivity reactions to any of its components. **WARNINGS:** Because of the potential hazard of nephrotoxicity due to the polyethylene glycol content of the base, care should be exercised when using this product in treating extensive deep lesions where absorption of large quantities of polyethylene glycol is possible. Safety of use in pregnant or breeding animals has not been determined. MURICIN Ointment is not for ophthalmic use. **ADVERSE REACTIONS:** No adverse reactions have been reported with this product. If a skin reaction such as irritation should occur, treatment should be discontinued and appropriate therapy instituted. **DOSAGE AND ADMINISTRATION:** Prior to treatment, the lesion should be cleansed. MURICIN Ointment should be applied to the affected area twice a day. Apply a sufficient amount of ointment to completely cover the infected area. Maximum duration of treatment should not exceed 30 days. **HOW SUPPLIED:** MURICIN Ointment is supplied in 15-gram tubes, NDC 17033-420-15. Store between 15° and 30°C (59° and 86°F). Keep out of reach of children.




1. Remove entire Applicator Assembly, puncture tube with white portion of the cap.



2. Replace entire Applicator Assembly.



3. To dispense, remove the white portion of the cap.



4. After use replace white cap to close.

IMPORTANT: The opening of this product is covered by a metal tamper-resistant seal. If this seal has been punctured or is not visible, do not use and return product to place of purchase.

Manufactured for:
Dechra Veterinary Products
Overland Park, KS 66211


Dechra
Veterinary Products